



Product Datasheet

Product Name	Angiopoietin Like Protein 3 Human Recombinant
Cata No	CB500150
Source	<i>Escherichia Coli.</i>
Synonyms	Angiopoietin 5, ANGPT5, ANGPTL3, Angiopoietin Like Protein 3.

Description

ANGPTL3 and ANGPTL4 are angiopoietin-like proteins secreted and expressed mainly by the liver, their role being the regulation of triglyceride metabolism by inhibiting the lipolysis of triglyceride-rich lipoproteins. During different nutritional states (feeding/fasting) the levels of the circulating triglycerides are regulated by Angptl3 and Angptl4 through differential inhibition of Lipoprotein lipase (LPL) as shown by the experimental data. The molecular structure of ANGPTL3 is similar to that of the angiopoietins (vascular endothelial growth factors). Deletion mutants of human Angiopoietin 5 were used in order to demonstrate that the N-terminal domain (fragment 17-207) and not the C-terminal fibrinogen-like domain (fragment 207-460) increased the plasma triglyceride levels in mice. The ANGPTL3 Human Recombinant is produced with N-terminal fusion of His-Tag.

The Angiopoietin-like protein 3 His Tagged Fusion Protein is 26kDa containing 207 amino acid residues of the ANGPTL3 Human and 16 additional amino acid residues – His-Tag (underlined).

MRGSHHHHHH GMASHMSRID QDNSSFDLSL
PEPKSRFAML DDVKILANGL LQLGHGLKDF
VHKTKGQIND IFQKLNIFDQ SFYDLSLQTS
EIKEEEKELR RTTYKLQVKN EEVKNMSLEL
NSKLESLL EE KILLQQVKY LEEQLTNLIQ
NQPETPEHPE VTSLKTFVEK QDNSIKDLLQ

TVEDQYKQLN QQHSQIKEIE NQLRRTSIQE
PTEISLSSKP RAP.

Purity

Angiopoietin 5 purity is greater than 95% as determined by SDS-PAGE.

Formulation

ANGPTL3 Human Sterile filtered and lyophilized from 0.5 mg/ml in 0.05M Acetate buffer pH-4.

Solubility

Add 0.2 ml of 0.1M Acetate buffer pH-4 and let the lyophilized pellet of ANGPTL3 Human dissolve completely. For conversion into higher pH value, we recommend intensive dilution by relevant buffer to a concentration of 10µg/ml. In higher concentrations the solubility of Angiopoietin 5 is limited.

Stability

Store lyophilized ANGPTL3 Human at -20°C. Aliquot the product after reconstitution to **avoid repeated freezing/thawing cycles**. Reconstituted Angiopoietin 5 can be stored at 4°C for a limited period of time; it does not show any change after two weeks at 4°C.

The lyophilized Angiopoietin 5 remains stable until the expiry date when stored at -20°C.

Application

Western blotting, ELISA.

*** For Non-Clinical Research Use Only ***